

Table of Contents

Post processing tools.....	1
Standard Post-Processing.....	1
Calculation of Residual Circulation and EP-fluxes.....	1
Required Input.....	1
EPF-Routines.....	1
Other Useful Scripts.....	1
CMOR (Climate Model Output Rewriter).....	2

Post processing tools

Standard Post-Processing

CDO is a very useful tool for postprocessing SOCOL output, e.g. for calculating averages, interpolation to pressure levels, selection of variables, time steps, etc.: <https://code.mpimet.mpg.de/projects/cdo/>

- cdo is already installed on the IAC Linux system and can be loaded by `module load cdo`

The classical post processor for ECHAM data is the "afterburner" which provides the following operations:

- Extract specified variables and levels
- Compute derived variables
- Transform spectral data to Gaussian grid representation
- Vertical interpolation to pressure levels or height
- Write data in GRIB, NetCDF, SERVICE or EXTRA format
- User manual: afterburner.txt
- Afterburner source code and Makefile: after.tar
- ⚠ The cdo command *after* does the same as the afterburner, and is easier to use!!!

Calculation of Residual Circulation and EP-fluxes

The calculation of the residual circulation and the EP-fluxes follow Andrews (1987):
https://wiki.iac.ethz.ch/pub/Chemie/PostProc/Andrews_1987_p127ff.pdf

Required Input

- daily mean or 6 hrly (better) instantaneous 3D T [K], u [m/s], v [m/s], omega [Pa/s] on pressure levels [Pa]
- ⚠ ATTENTION: the namelist setting `PUTDATA = 24, 'hours', 'first', 0` outputs **instantaneous values every 24 hours, not daily averages!!!** That leads to artefacts in the calculated TEM in the mesosphere. If you intend to calculate the residual circulation from your model output, set `PUTDATA = 12, 'hours', 'first', 0` or better `PUTDATA = 6, 'hours', 'first', 0`
- cdo script for preparing input data for epf routine from model output:
https://wiki.iac.ethz.ch/pub/Chemie/PostProc/make_epf_vars_netcdf.sh

EPF-Routines

- Using a temperature-dependent density:
https://wiki.iac.ethz.ch/pub/Chemie/PostProc/epf_from_daily_v3.f90
- Using a fixed scale height of $H=6950\text{ m}$ and log-pressure altitude:
https://wiki.iac.ethz.ch/pub/Chemie/PostProc/epf_from_daily_v4.f90
- ⚠ NOTE: Although CCMI recommended a fixed scale height, we used a temperature-dependent density to calculate the residual circulation and EP-fluxes from the CCMI simulations!!!

Other Useful Scripts

- Compiling epf routines: https://wiki.iac.ethz.ch/pub/Chemie/PostProc/compile_epf_from_daily.sh
- Script to postprocess several months: https://wiki.iac.ethz.ch/pub/Chemie/PostProc/calc_ep-flux.csh

CMOR (Climate Model Output Rewriter)

CMOR is a "handy" data post-processing tool originally designed to produce datasets conforming to CMIP conventions. It has since been adapted by the CCMi community for CCMi-1. CMOR stands for Climate Model Output Rewriter.

- Laura's CMOR documentation:
https://wiki.iac.ethz.ch/pub/Chemie/PostProc/CMOR_SOCOL_documentation.pdf
- Laura's CMOR scripts: https://wiki.iac.ethz.ch/pub/Chemie/PostProc/cmor_scripts.tar

Edit | Attach | Print version | History: %REVISIONS% | Backlinks | Raw View | More topic actions
Topic revision: r6 - 12 Apr 2022 - 17:59:48 - AndreaStenke

- ☐ Chemie

No permission to view Chemie.WebLeftBar

•
•



Copyright © by the contributing authors. All material on this collaboration platform is the property of the contributing authors.

Ideas, requests, problems regarding Wiki? Send feedback